

### 3.6 Properties of Rational Functions

A **rational function** is a function of the form  $R(x) = \frac{p(x)}{q(x)}$ , where  $p$  and  $q$  are polynomial functions and  $q$  is not the zero polynomial.

#### Finding the Domain of a Rational Function

The domain of a rational function is the set of all real numbers except those for which the denominator  $q$  is 0.

**Examples:** Find the domain of each rational function.

$$\text{a) } R(x) = \frac{2x^2 - 3}{x + 3}$$

$$\text{b) } R(x) = \frac{x + 3}{x^2 - 9}$$

$$\text{c) } R(x) = \frac{x^2}{x^2 + 7x + 12}$$

★ **Note:** It is important to understand that  $R(x) = \frac{x + 3}{x^2 - 9}$  and  $f(x) = \frac{1}{x - 3}$  are not the same. They have different domains. Their graphs are nearly identical, but the graph of the first function has a hole in it at  $x = -3$ , while the graph of the second function does not.

A rational function  $R(x) = \frac{p(x)}{q(x)}$  is in **lowest terms** if  $p(x)$  and  $q(x)$  have no common factors.

#### Finding the Intercepts of a Rational Function

★ **Warning:** Always write the rational function in lowest terms before finding the  $x$ -intercepts of the graph. Otherwise, you may end up listing values that are actually holes in the graph. When finding the  $y$ -intercepts, remember that if zero is not in the domain, there is no  $y$ -intercept.

To find the  $x$ -intercepts (real zeros) of the graph of a rational function, we set  $R(x) = 0$  and solve for  $x$ .

Notice that if  $R(x) = \frac{p(x)}{q(x)} = 0$ ,  $p(x)$  must equal zero. It is the numerator that tells us about the  $x$ -intercepts.

- To find the  **$x$ -intercepts** (real zeros), simplify the rational function and set the numerator equal to zero.
- To find the  **$y$ -intercept**, plug  $x = 0$  into either the simplified or the unsimplified function. If zero is not in the domain of the function, there is no  $y$ -intercept.


**Examples:** Find the  $x$ - and  $y$ - intercepts of each rational function.

$$\text{a) } R(x) = \frac{x + 2}{x^2 - 16}$$


$$\text{b) } R(x) = \frac{x^2 - 2x - 35}{x^2 + 11x + 30}$$

$$\text{c) } R(x) = \frac{x}{x^2 + 8x}$$

**Example:** Analyze the graph of  $H(x) = \frac{1}{x^2}$ . (Find the domain, range, intercepts, and any other important features).



**Example:** Using transformations, graph  $R(x) = \frac{1}{(x+2)^2} - 1$ .



### Asymptotes

In the previous example, notice that as the values of  $x$  become more negative, that is, as  $x$  becomes **unbounded in the negative direction** ( $x \rightarrow -\infty$ , read "as  $x$  approaches negative infinity") the values of  $R(x)$  approach  $-1$ .





We conclude the following:

1. As  $x \rightarrow -\infty$ , the values of  $R(x) \rightarrow -1$ .
2. As  $x \rightarrow -2$ , the values of  $R(x) \rightarrow \infty$ .
3. As  $x \rightarrow \infty$ , the values of  $R(x) \rightarrow -1$ .

The graph goes near, but does not reach the lines  $x = -2$  and  $y = -1$ . These lines are called **asymptotes**.

Let  $R$  denote a function.

- If, as  $x \rightarrow -\infty$  or as  $x \rightarrow \infty$ , the values of  $R(x)$  approach some fixed number  $L$ , then the line  $y = L$  is a **horizontal asymptote** of the graph of  $R$ .
- If, as  $x$  approaches some number  $c$ , the values of  $R(x)$  approach  $\infty$  or  $-\infty$ , then the line  $x = c$  is a **vertical asymptote** of the graph of  $R$ .



There is a third type of asymptote called an **oblique asymptote**. An oblique asymptote occurs when the graph's end behavior follows an oblique line (a linear function of the form  $y = ax + b$ ,  $a \neq 0$ ). A graph may also approach some function such as a parabola.

★ **Note:** The graph of a function may intersect a horizontal or oblique asymptote at some other point, but the graph will never intersect a vertical asymptote.

### Find the Vertical Asymptotes of a Rational Function

A rational function  $R(x) = \frac{p(x)}{q(x)}$ , in *lowest terms* will have a vertical asymptote at  $x = r$  if  $r$  is a real zero of the denominator  $q$ . That is, if  $x - r$  is a factor of the denominator  $q$  of the rational function in lowest terms, it will have the vertical asymptote  $x = r$ .

- **Warning:** If the rational function is not in lowest terms, this theorem will result in an incorrect listing of vertical asymptotes.

**Example:** Find the vertical asymptotes, if any, of the graph of each rational function.

a)  $R(x) = \frac{x+1}{x^2-9}$

b)  $R(x) = \frac{x+2}{x^2-3x-10}$

c)  $R(x) = \frac{x}{x^3+10x^2+9x}$

d)  $R(x) = \frac{x^2}{x^2+4}$

### Find the Horizontal or Oblique Asymptotes of a Rational Function

- If a rational function  $R(x)$  is **proper**, that is *if the degree of the numerator is less than the degree of the denominator*, then the line  $y = 0$  is a horizontal asymptote.
- If a rational function  $R(x) = \frac{p(x)}{q(x)}$  is **improper**, that is, *if the degree of the numerator is greater than or equal to the degree of the denominator*, we can use long division to write the rational function as the sum of a polynomial  $f(x)$  plus the proper rational function  $\frac{r(x)}{q(x)}$ . As  $x \rightarrow -\infty$  or as  $x \rightarrow \infty$ ,  $\frac{r(x)}{q(x)} \rightarrow 0$ , and the graph of  $R(x)$  approaches the graph of  $f(x)$ .
  - The possibilities include:
    1. If  $f(x) = b$ , a constant, the line  $y = b$  is a horizontal asymptote of the graph of  $R$ .
    2. If  $f(x) = ax + b$ ,  $a \neq 0$ , the line  $y = ax + b$  is an oblique (slanted) asymptote of the graph of  $R$ .
    3. In all other cases, the graph of  $R$  approaches the graph of  $f$ , and there are no horizontal or oblique asymptotes.

**Examples:** Find the horizontal or oblique asymptotes, if any, of the graph of the function.

a)  $R(x) = \frac{x+5}{x^2-3x-10}$

b)  $H(x) = \frac{6x^2-2x+5}{3x^2-2}$

c)  $H(x) = \frac{8x^3+2x^2-6}{2x^2-3}$

d)  $H(x) = \frac{3x^4+4x^2-7}{x^2-2x+5}$

## Summary:

Given a rational function  $R(x) = \frac{p(x)}{q(x)}$ ,

1. The **domain** is found by setting the denominator not equal to zero and solving. (Do not simplify first).
2. The **x-intercepts** are found by setting the numerator to zero and solving. (Simplify first).
3. The **y-intercept** is found by setting  $x = 0$  and simplifying. (You can do this before or after simplifying the function). Remember, if zero is not in the domain, the function has no y-intercept.
4. **Vertical asymptotes** are found by setting the denominator to zero and solving (Simplify first).
5. **Horizontal or Oblique Asymptotes:**
  - a) If **proper**, where *degree of numerator* < *degree of denominator*, the graph has a **horizontal asymptote** at  $y = 0$ .
  - b) If **improper**, where *degree of numerator* = *degree of denominator*, the graph has a **horizontal asymptote** at the line  $y = \frac{\text{leading coefficient of the numerator}}{\text{leading coefficient of the denominator}}$ .
  - c) If **improper**, where *degree of numerator* > *degree of denominator (by one)*, the graph has an **oblique asymptote** of  $y = f(x)$  found by performing long division.
  - d) If **improper**, where *degree of numerator* > *degree of denominator (by more than one)*, then  $R$  has neither a horizontal nor an oblique asymptote, but the end behavior can be determined using long division.